arXiv:1304.3392v1 [math.CA] 11 Apr 2013

LOCALIZATION AND DIMENSION FREE ESTIMATES FOR MAXIMAL FUNCTIONS.

ALBERTO CRIADO AND FERNANDO SORIA

ABSTRACT. In the recent paper [J. Funct. Anal. **259** (2010)], Naor and Tao introduce a new class of measures with a so-called micro-doubling property and present, via martingale theory and probability methods, a localization theorem for the associated maximal functions. As a consequence they obtain a weak type estimate in a general abstract setting for these maximal functions that is reminiscent of the '$n \log n$ result' of Stein and Strömberg in Euclidean spaces.

The purpose of this work is twofold. First we introduce a new localization principle that localizes not only in the time-dilation parameter but also in space. The proof uses standard covering lemmas and selection processes. Second, we show that a uniform condition for micro-doubling in the Euclidean spaces provides indeed dimension free estimates for their maximal functions in all L^p with $p > 1$. This is done introducing a new technique that allows to differentiate through dimensions.

1. INTRODUCTION

We say that (X, d, μ) is a metric measure space if (X, d) is a separable metric space and μ a Radon measure on it. We denote by $B(x, r)$ the open ball centered at x with radius r with respect to the metric d, that is

$$B(x,r) = \{y \in X : d(x,y) < r\}.$$

We will assume that the measure μ is non-degenerate. This means that any ball with positive radius has non-zero measure. Given $T \subset (0,\infty)$, for a locally integrable function f over X we define the following centered maximal operator

$$M_T f(x) = \sup_{r \in T} \frac{1}{\mu(B(x,r))} \int_{B(x,r)} |f(y)| \, d\mu(y).$$

When $T = (0,\infty)$, the operator $M = M_T$ represents the usual Hardy-Littlewood maximal operator. As in the case of evolution equations and semigroup theory we can think of T as a set of times and then M_T is the maximal operator on them.

We will also assume that the measure μ satisfies a doubling property. This simply says that the measure of a ball is comparable with the measure of certain dilation of it. The classical way of expressing the doubling property uses the dilation factor 2. That is, μ is doubling if there exists a constant $K > 0$ so that for each $x \in X$ and $R > 0$ one has $\mu(B(x, 2R)) \leq K\mu(B(x, R))$.

In these hypotheses one can reproduce the argument of Vitali's covering lemma to show that M_T is weakly bounded on $L^1(\mu)$. That is, there exist a constant $c_{\mu,1} > 0$ so that

$$\mu(\{x \in X : M_T f(x) > \lambda\}) \leq \frac{c_{\mu,1}}{\lambda} \int_X |f| d\mu,$$

for all $\lambda > 0$ and all locally integrable f over X. Since $\|M_T f\|_{L^\infty(\mu)} \leq \|f\|_{L^\infty(\mu)}$, by interpolation one obtains the $L^p(\mu)$ bounds

$$\|M_T f\|_{L^p(\mu)} \leq C_{\mu,p} \|f\|_{L^p(\mu)},$$

2010 *Mathematics Subject Classification.* 42B25.

Key words and phrases. Maximal functions, radial measures, dimension free estimates.

Authors supported by DGU grant MTM2010-16518.

for all $p > 1$. The order of magnitude of the constants $c_{\mu,1}$ and $C_{\mu,p}$ that we obtain is essentially that of the doubling constant K and $K^{1/p}p/(p-1)$ respectively.

In the case that $X = \mathbb{R}^n$ and $\mu = m_n$ is Lebesgue measure, the value of the doubling constant K is 2^n. Hence the aforementioned constants $c_{m_n,1}$ and $C_{m_n,p}$ grow exponentially to infinity with the dimension. It has been a matter of interest to know if these constants can be bounded uniformly in dimension. E.M. Stein [31] (detailed proof in [32]) proved this for $C_{m_n,p}$ with $p > 1$ when considering the Euclidean metric. J. Bourgain [7], [8], [9] and independently A. Carbery [12], showed that for the metric given by a norm, uniform bounds hold if $p > 3/2$. This was extended to all the range $p > 1$ by D. Müller in [27] for the maximal functions associated with the ℓ^q metrics, with $1 \leq q < \infty$. Observe that this excludes the case $q = \infty$, where the 'balls' are cubes with sides parallel to the coordinate axes. Recently J. Bourgain has solved this case by giving uniform bounds on L^p for all $p > 1$ (see [11]).

The case $p = 1$ is more complicated. J.M. Aldaz showed in [2] that for the ℓ^∞ metric the constants $c_{m_n,1}$ grow to infinity as $n \to \infty$. As shown by Aubrun [6], in this case one has the estimate $c_{m_n,1} \geq C_\varepsilon (\log n)^{1-\varepsilon}$ for each $\varepsilon > 0$. It is still unknown if this also happens for other metrics, although there are partial results in the form of upper bounds for the possible growth of the constants. When considering the Euclidean metric a special argument allows to prove that $c_{m_n,1} = \mathcal{O}(n)$. For metrics given by general norms the best upper bound remains $c_{m_n,1} = \mathcal{O}(n \log n)$. Both results are due to E.M. Stein and J.O. Strömberg in [32]. In this last paper, instead of the usual doubling condition, they used the fact that in $\mathbb{R}^n$ one has

$$m_n(B(x,(1+1/n)R)) = (1+1/n)^n \, m_n(B(x,R)) \leq e \, m_n(B(x,R)).$$

That is, the dilation by the factor $(1+1/n)$ does not increase essentially the volume of a ball. This allows to perform a more efficient covering argument with less overlapping than in Vitali's lemma.

A. Naor and T. Tao observed in [28] that in order to extend Stein and Strömberg '$n \log n$' bound to general metric measure spaces suffices to assume that dilations by the factor $(1+1/n)$ preserve essentially the volumes of balls and that the measures of intersecting balls with the same radius are comparable. We next explain this conditions and state their result in detail.

Let (X, d, μ) be a metric measure space. Given a positive number $n \geq 1$, we will say that μ is n-micro-doubling if there exists a constant $K_0 > 0$ such that for each $x \in X$ and $R > 0$ one has

$$\mu(B((x,(1+1/n)R)) \leq K_0 \, \mu(B(x,R)).$$

Of course n here is not necessarily related to any notion of dimension. We will refer to K_0 as the n-micro-doubling constant.

We will say that a measure μ is weakly-doubling if the measure of two intersecting balls with the same radius is comparable. That is, there exists a constant $K_1 > 0$ such that for each $x, y \in X$ and $R > 0$ such that $B(x,R) \cap B(y,R) \neq \emptyset$ one has

$$\mu(B(y,R)) \leq K_1 \, \mu(B(x,R)).$$

Following the terminology of A. Naor and T. Tao in [28], we will say that a measure μ is strong n-micro-doubling if it is both, n-micro-doubling and weakly-doubling. Equivalently, μ is strong n-micro-doubling if there exist a constant $K > 0$ so that for each $x \in X$, $R > 0$, and $y \in B(x,R)$ one has

$$\mu(B(y,(1+1/n)R)) \leq K \, \mu(B(x,R)).$$

It is easy to see that every Ahlfors-David n-regular metric measure space is strong n-micro-doubling. We recall that a metric measure space (X, d, μ) is said to be Ahlfors-David n-regular if there exists a constant $C \geq 1$ so that

$$\frac{1}{C}\, R^n \leq \mu(B(x, R) \leq C\, R^n,$$

for all $x \in X$ and $R > 0$. In the Euclidean n-dimensional case, some examples of strong n-micro-doubling measures are given by the power densities $|x|^\alpha$, with $\alpha > -n$. See Section 3.1 for more details.

We will consider maximal operators associated with the following type of time sequences in $(0, \infty)$. A sequence $\{a_k\}_{k\in\mathbb{Z}} \subset (0, \infty)$ is lacunary if there exist $a > 1$ so that $a_{k+1} > aa_k$. If $a = n > 1$ in this setting of n-micro-doubling measures, we will say explicitly that the time sequence is n-lacunary. Now we are ready to state the maximal theorem presented by A. Naor and T. Tao in [28].

Theorem 1.1. *Let (X, d, μ) be a measure metric space, with μ a strong n-micro-doubling measure with constant K. Then we have the following weak type estimates*

i) if $T \subset (0, \infty)$ is an n-lacunary sequence, then [1]

$$\mu(\{x \in X : M_T f(x) > \lambda\}) \leq \frac{C_K}{\lambda} \|f\|_{L^1(\mu)},$$

ii) if $T \subset (0, \infty)$ is a lacunary sequence with constant a, then

$$\mu(\{x \in X : M_T f(x) > \lambda\}) \leq \frac{C_K}{\lambda} \frac{\log n}{\log a} \|f\|_{L^1(\mu)},$$

iii) if $T = (0, \infty)$ so that $M = M_T$ is the Hardy-Littlewood maximal operator, then

$$\mu(\{x \in X : Mf(x) > \lambda\}) \leq \frac{C_K n \log n}{\lambda} \|f\|_{L^1(\mu)},$$

where the C_K are constants only depending on K.

Note that part $iii)$ is a generalization of the '$n \log n$' bound by E.M. Stein and J.O. Strömberg (see [32]) in normed Euclidean spaces. In this setting, parts $i)$ and $ii)$ are due to M.T. Menárguez and the second author in [25], as well as the chain of implications $i) \Rightarrow ii) \Rightarrow iii)$.

Moreover, A. Naor and T. Tao showed also in [28] that the $\mathcal{O}(n \log n)$ bound is optimal even in the setting of Ahlfors-David n-regular spaces, by constructing a sequence of such spaces (X_n, d_n, μ_n) so that

$$\|M_{\mu_n}\|_{L^1(\mu_n) \to L^{1,\infty}(\mu_n)} \geq Cn \log n.$$

Theorem 1.1 can be proved as in [25] or by an argument of Lindenstrauss present in [22] (see also [28]). However, the main contribution of A. Naor and T. Tao in [28] is the following nice localizaton principle for maximal operators, from which they obtained Theorem 1.1 as a corollary.

---

[1] Statement $i)$ of the theorem does not appear explicitly in [28].

Theorem 1.2. *Let (X, d, μ) be a metric measure space and let μ be n-micro-doubling with constant K_0. If $T \subset (0, \infty)$ and $p \geq 1$, then we have the following localizaton property for the weak L^p norms of the associated maximal operator*

$$\|M_T\|_{L^p \to L^{p,\infty}} \leq C_{K_0} + C'_{K_0} \sup_{k \in \mathbb{Z}} \|M_k\|_{L^p \to L^{p,\infty}}.$$

Here M_k denotes the restriction of M_T to times in $[n^k, n^{k+1}]$, that is $M_k = M_{T \cap [n^k, n^{k+1}]}$, and C_{K_0} and C'_{K_0} are constants only depending on K_0.[2]

The proof given by A. Naor and T. Tao of this result is probabilistic and relies on random martingales and Doob-type maximal inequalities. Here we present (see Theorem 2.1) a more geometrical proof based on covering lemmas and selection processes that is closer in spirit to the arguments in [32]. In addition, our result considers localization not only in time but also in space (see the statement of the theorem). The proof of this and the connection with Theorem 1.1 will be presented in Section 2.

In the rest of the paper we restrict ourselves to the Euclidean case. We study measures given by a radial density, so that they can be defined in all dimensions. It is known that if these measures are finite, and hence are neither weakly-doubling nor n-micro-doubling, the (weak) L^p operator norms of the associated maximal functions grow exponentially to infinity with the dimension at least for a range of values of p near 1 (see [1], [3] and [13]). There are cases in which this range of p's may consist of the whole interval $[1, \infty)$. An example of this is given by the Gaussian measure (see [15]).

The situation changes if we consider measures that are strong n-micro-doubling in each $\mathbb{R}^n$ with constants uniformly bounded in dimension. In this case Theorem 1.1 applies and the weak L^1 operator norms of the maximal function grow at most like $\mathcal{O}(n \log n)$. In Section 3.1 we study maximal operators associated with uniformly weakly doubling measures. For such measures μ the main result of this section asserts that M_μ is uniformly bounded on $L^p(\mu, \mathbb{R}^n)$ for all $p > 1$. The same thing is true in weak L^1 but restricting the action of M_μ to radial functions.

The previous results are obtained via the following characterization: μ is uniformly weakly doubling if and only if its density is essentially constant over dyadic annuli. This equivalence is proved using a new method of differentiation through dimensions presented in Section 3.2.

The case of decreasing densities is treated in Section 3.4. The situation here is that the uniform weakly doubling property can be obtained from surprisingly mild conditions.

Finally, Section 4 contains examples of doubling measures with associated maximal operators failing to have uniform bounds.

2. LOCALIZATION PRINCIPLES

In this section we will obtain Theorems 1.1 and 1.2 as a consequence of yet another localization theorem, whose proof avoids technical arguments from probability theory. The statement is the following:

---

[2] The constants C_{K_0} and C'_{K_0} obtained in [28] are of the order of $\mathcal{O}(K_0)$ and $\mathcal{O}(1+\log\log K_0/(1+\log n))$ respectively.

Theorem 2.1. *Let (X, d, μ) be a metric measure space and let μ be n-micro-doubling with constant K_0. If $T \subset (0, \infty)$ and $p \geq 1$ are fixed, then for each locally integrable function f over X and each $\lambda > 0$ one can find a disjoint collection of measurable sets $\{A_k\}_{k\in\mathbb{Z}}$ such that*

$$\begin{aligned} &\mu(\{x \in X : \lambda < M_T f(x) \leq 2\lambda\}) \\ &\qquad \leq \quad C_1 \left(\frac{1}{\lambda^p} \|f\|^p_{L^p(\mu)} + \sum_{k\in\mathbb{Z}} \mu(\{x \in X : M_k(f\chi_{A_k})(x) > C_2\lambda\}) \right), \end{aligned}$$

where C_1 and C_2 are constants that only depend on K_0.

It is not difficult to show that Theorem 2.1 implies Theorem 1.2. To see this, given a locally integrable f and $\lambda > 0$ we write

$$\begin{aligned} E_\lambda &= \{x \in X : M_T f(x) > \lambda\}, \\ F_\lambda &= \{x \in X : \lambda < M_T f(x) \leq 2\lambda\}. \end{aligned}$$

Note that by the disjointness of the collection $\{A_k\}_{k\in\mathbb{Z}}$ we have

$$\begin{aligned} \sum_{k\in\mathbb{Z}} \mu(\{x \in X : M_k(f\chi_{A_k})(x) > C_2\lambda\}) &\leq \frac{C_2^{-p}}{\lambda^p} \sum_{k\in\mathbb{Z}} \|M_k\|^p_{L^p\to L^{p,\infty}} \int_{A_k} |f|^p \, d\mu \\ &\leq \frac{C_2^{-p}}{\lambda^p} \sup_{k\in\mathbb{Z}} \|M_k\|^p_{L^p\to L^{p,\infty}} \int_X |f|^p \, d\mu. \end{aligned}$$

Hence Theorem 2.1 gives the estimate

$$\mu(F_\lambda) \leq \frac{C_{K_0}}{\lambda^p} \left(1 + \sup_{k\in\mathbb{Z}} \|M_k\|^p_{L^p\to L^{p,\infty}}\right) \|f\|^p_{L^p(\mu)}.$$

This implies

$$\begin{aligned} \mu(E_\lambda) &= \mu\left(\bigcup_{j=0}^{\infty} F_{2^j\lambda}\right) = \sum_{j=0}^{\infty} \mu(F_{2^j\lambda}) \\ &\leq \sum_{j=0}^{\infty} \frac{C_{K_0}}{(2^j\lambda)^p} \left(1 + \sup_{k\in\mathbb{Z}} \|M_k\|^p_{L^p\to L^{p,\infty}}\right) \|f\|^p_{L^p(\mu)} \\ &= \frac{C_{K_0}}{\lambda^p} \left(1 + \sup_{k\in\mathbb{Z}} \|M_k\|^p_{L^p\to L^{p,\infty}}\right) \|f\|^p_{L^p(\mu)}, \end{aligned}$$

as wanted.

Also part $i)$ of Theorem 1.1 follows from Theorem 1.2 using the following argument.

If $T = \{a_j\}_{j\in\mathbb{Z}}$ is an n-lacunary sequence then each interval of the form $[n^k, n^{k+1}]$ contains at most one element of T. Theorem 1.2 implies that

$$\|M_T\|_{L^1\to L^{1,\infty}} \leq C_K \sup_{j\in\mathbb{Z}} \|M_{\{a_j\}}\|_{L^1\to L^{1,\infty}}.$$

We only have to show now that $M_{\{a_j\}}$ is bounded independently of j. Since

$$M_{\{a_j\}} f(x) = \frac{1}{\mu(B(x, a_j))} \int_{B(x,a_j)} |f(y)| \, d\mu(y),$$

by the weak doubling property of μ and Fubini Theorem we have

$$\begin{aligned}\|M_{\{a_j\}}f\|_{L^1} &= \int_X \frac{1}{\mu(B(x,a_j))}\int_X \chi_{B(x,a_j)}(y)|f(y)|\,d\mu(y)\,d\mu(x)\\ &\le \int_X\int_X \frac{K_1}{\mu(B(y,a_j))}\chi_{B(y,a_j)}(x)|f(y)|\,d\mu(y)\,d\mu(x)\\ &= \int_X \frac{K_1}{\mu(B(y,a_j))}\int_{B(y,a_j)} d\mu(x)\,|f(y)|\,d\mu(y) = K_1\|f\|_{L^1}.\end{aligned}$$

This says, in particular, that each $M_{\{a_j\}}$ is bounded on L^1 with operator norm bounded by K_1.

The implications $i) \Rightarrow ii)$ and $ii) \Rightarrow iii)$ of Theorem 1.1 can be obtained through the same generic arguments given in [25]. The first one says that every maximal operator associated with a lacunary sequence is essentially majorized by $\log n$ operators associated with n-lacunary sequences. The second one says that the full maximal operator is majorized by n operators, each associated with a lacunary sequence.

We now give a proof of Theorem 2.1.

Proof of Theorem 2.1. Given a locally integrable f and $\lambda > 0$ we write

$$F_\lambda = \{x \in X : \lambda < M_T f(x) \le 2\lambda\}.$$

Once we determine what the sets A_j are, we will have to prove that

$$\mu(F_\lambda) \le C_1 V^f_{\lambda,p},$$

where we have used the notation

$$V^f_{\lambda,p} = \frac{1}{\lambda^p}\|f\|^p_{L^p(\mu)} + \sum_{k\in\mathbb{Z}} \mu(\{x \in X : M_k(f\chi_{A_k})(x) > C_2\lambda\}).$$

Note first that it is enough to prove the result for $p = 1$. To see this, given a function f in $L^p(\mu)$ we consider $f = f^\lambda + f_\lambda$ where

$$\begin{aligned} f^\lambda &= f\chi_{\{|f|>\lambda\}} \in L^1(\mu),\\ f_\lambda &= f\chi_{\{|f|\le\lambda\}} \in L^\infty(\mu).\end{aligned}$$

Since

$$M_T f(x) \le M_T f^\lambda(x) + M_T f_\lambda(x) \le M_T f^\lambda(x) + \lambda,$$

for $M_T f(x) > 2\lambda$ to hold it is necessary that $M_T f^\lambda(x) > \lambda$. Hence,

$$F_{2\lambda} \subset \{x \in X : \lambda < M_T f^\lambda(x) \le 4\lambda\} = G_1 \cup G_2,$$

with

$$\begin{aligned} G_1 &= \{x \in X : \lambda < M_T f^\lambda(x) \le 2\lambda\},\\ G_2 &= \{x \in X : 2\lambda < M_T f^\lambda(x) \le 4\lambda\}.\end{aligned}$$

Then since $f^\lambda \in L^1(\mu)$ we apply the result for $p = 1$ to G_1 and G_2 to obtain

$$\begin{aligned}\mu(F_{2\lambda}) &\le 2\max(\mu(G_1),\mu(G_2))\\ &\le C\left(C_1\int_X \frac{|f^\lambda|}{\lambda}\,d\mu + \sum_{k\in\mathbb{Z}}\mu(\{x\in X : M_k(f^\lambda\chi_{A_k})(x) > C_2\lambda\})\right)\\ &\le C\left(C_1\int_X \frac{|f|^p}{\lambda^p}\,d\mu + \sum_{k\in\mathbb{Z}}\mu(\{x\in X : M_k(f\chi_{A_k})(x) > C_2\lambda\})\right),\end{aligned}$$

for each $p \geq 1$. Since λ is arbitrary the result is proved for any $p \geq 1$.

Now we prove the result in the case that $p = 1$, that is

$$\mu(F_\lambda) \leq C_1 V^f_{\lambda,1}$$

We consider the following collections of balls

$$\begin{aligned} \mathcal{B} &= \{B(x,R) : x \in X,\ R \in T\}, \\ \mathcal{B}_k &= \{B(x,R) : x \in X,\ R \in T \cap [n^k, n^{k+1}]\}. \end{aligned}$$

Given a ball $B \in \mathcal{B}$ we denote by z_B its center and by R_B its radius. We define the collection

$$\mathcal{A} = \left\{ B \in \mathcal{B} : \lambda < \frac{1}{\mu(B)} \int_B |f| \, d\mu \leq 2\lambda \right\}.$$

For each $B \in \mathcal{A}$ one can find a concentric ball $\tilde{B}$ so that $R_B/(1+1/n) \leq R_{\tilde{B}} < R_B$ and

$$\frac{1}{\mu(B)} \int_{\tilde{B}} |f| \, d\mu > \lambda.$$

We will write $B^0 = B(z_B, R_B - R_{\tilde{B}})$. Since

$$F_\lambda \subset \bigcup_{B \in \mathcal{A}} B^0,$$

it suffices to prove that

$$\mu\left(\bigcup_{B \in \mathcal{A}} B^0 \right) \leq C_1 V^f_{\lambda,1}.$$

It is not difficult to see (see the argument preceding (2.4) below) that $\cup_{B \in \mathcal{A}} B^0$ is contained in a level set of $M_T f$ and, as a consequence, has finite μ-measure. Using that X is separable and the monotone convergence there exists $\mathcal{A}' \subset \mathcal{A}$ with $\sharp \mathcal{A}' < \infty$ such that

$$\mu\left(\bigcup_{B \in \mathcal{A}'} B^0 \right) \geq \frac{1}{2} \mu\left(\bigcup_{B \in \mathcal{A}} B^0 \right).$$

Hence, it is enough to show that

$$\mu\left(\bigcup_{B \in \mathcal{A}'} B^0 \right) \leq C_1 V^f_{\lambda,1},$$

with C independent of $\mathcal{A}'$. Writing

$$\mathcal{A}_k := \{B \in \mathcal{A}' : B \in \mathcal{B}_k\},$$

we have that $\mathcal{A}' = \mathcal{A}^1 \cup \mathcal{A}^2$ with

$$\begin{aligned} \mathcal{A}^1 &= \bigcup_{k \text{ odd}} \mathcal{A}_k, \\ \mathcal{A}^2 &= \bigcup_{k \text{ even}} \mathcal{A}_k. \end{aligned}$$

Assume that $\mu(\bigcup_{B \in \mathcal{A}^1} B^0) \geq \mu(\bigcup_{B \in \mathcal{A}^2} B^0)$ (if not, interchange the names), then $\mu(\bigcup_{B \in \mathcal{A}^1} B^0) \geq \frac{1}{2}\mu(\bigcup_{B \in \mathcal{A}'} B^0)$ and we just need to prove that

$$\mu\left(\bigcup_{B \in \mathcal{A}^1} B^0 \right) \leq C_1 V^f_{\lambda,1}.$$

For the sake of simplicity in the notation we rename $\mathcal{A} = \mathcal{A}^1$.

Now we want to write the set $\bigcup_{B\in\mathcal{A}} B^0$ as a union of disjoint sets. Observe that $\mathcal{A}=\{B_j\}_{j=1,\cdots,J}$ for certain J. We define $D_{B_1}=B_1^0$, and if $D_{B_1},\cdots,D_{B_m}$ are already defined, we take

$$D_{B_{m+1}}=B_{m+1}\setminus\bigcup_{j=1}^{m}B_j^0.$$

Then, we have

$$\mu\left(\bigcup_{B\in\mathcal{A}}B^0\right)=\mu\left(\bigcup_{B\in\mathcal{A}}D_B\right).$$

We also define the functions

$$g_B(x)=\frac{\mu(D_B)}{\mu(B)}\chi_{\tilde{B}}(x)$$

for $B\in\mathcal{A}$ and

$$G_k(x)=\sum_{B\in\mathcal{A}_k}g_B(x).$$

We start a selection process. Take k_1 as the largest $k\in\mathbb{Z}$ with $\mathcal{A}_k\neq\emptyset$, then we take $\tilde{G}_{k_1}=G_{k_1}$ and $\tilde{\mathcal{A}}_{k_1}=\mathcal{A}_{k_1}$. Once $\tilde{G}_{k_1},\cdots,\tilde{G}_{k_{m-1}}$ are determined for $m\geq 2$, we take k_m as the largest $k<k_{m-1}$ so that $\mathcal{A}_k\neq\emptyset$. We say that $B\in\tilde{\mathcal{A}}_{k_m}$ if $B\in\mathcal{A}_{k_m}$ and

$$\sum_{j=1}^{m-1}\tilde{G}_{k_j}\leq 1$$

on B. Then we define

$$\tilde{G}_{k_m}(x)=\sum_{B\in\tilde{\mathcal{A}}_{k_m}}g_B(x).$$

Since $\mathcal{A}$ is finite this process ends in a finite number of steps, and we have obtained $\tilde{G}_{k_1},\cdots,\tilde{G}_{k_M}$ for certain M. We call

$$\tilde{\mathcal{A}}=\bigcup_{m=1}^{M}\tilde{\mathcal{A}}_{k_m},$$

and claim that

$$\mu\left(\bigcup_{B\in\mathcal{A}}D_B\right)\leq(1+K_0)\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B). \tag{2.1}$$

To prove this claim note that if $A\in\mathcal{A}_{k_m}\setminus\tilde{\mathcal{A}}_{k_m}$, then there exists $z\in A$ such that

$$\sum_{j=1}^{m-1}\tilde{G}_{k_j}(z)>1.$$

Note that if $A\cap\tilde{B}\neq\emptyset$ for some $B\in\tilde{\mathcal{A}}_{k_j}$ with $j<m$, then $A\subset B^*:=B(x_B,(1+1/n)R_B)$. Thus for all $z\in A$ we have

$$\sum_{j=1}^{m-1}G^*_{k_j}(z):=\sum_{j=1}^{m-1}\sum_{B\in\tilde{\mathcal{A}}_{k_j}}\frac{\mu(D_B)}{\mu(B)}\chi_{B^*}(z)>1.$$

Then, by Tchebychev inequality

$$\begin{aligned}\mu\left(\bigcup_{B\in\mathcal{A}\setminus\tilde{\mathcal{A}}} D_B\right) &\le \mu\left(\Big\{z\in X : \sum_{j=1}^{M} G^*_{k_j}(z) > 1\Big\}\right)\\ &\le \sum_{j=1}^{M}\int_X G^*_{k_j}(z)\,d\mu(z) = \sum_{B\in\tilde{\mathcal{A}}}^{M}\frac{\mu(D_B)}{\mu(B)}\mu(B^*)\\ &\le K_0\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B).\end{aligned}$$

Now that the claim is justified, we only need to prove

$$\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B)\le C_1 V^f_{\lambda,1}. \tag{2.2}$$

By the definition of $\tilde{B}$ we have

$$\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B)\le\frac{1}{\lambda}\sum_{B\in\tilde{\mathcal{A}}}\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\,d\mu\,\mu(D_B)=\frac{1}{\lambda}\int_X|f|\left(\sum_{j=1}^{M}\tilde{G}_{k_j}\right)\,d\mu. \tag{2.3}$$

For each $j=1,\cdots,M$ we define $\tilde{A}_j := \operatorname{supp}\tilde{G}_{k_j}$. We now take $A_M=\tilde{A}_M$ and

$$A_j=\tilde{A}_j\setminus\bigcup_{\ell=j+1}^{M}\tilde{A}_\ell.$$

If $z\in A_m$ we have $\tilde{G}_{k_j}(z)=0$ if $j>m$, and then, by the way we selected $\tilde{G}_{k_m}$

$$\sum_{j=1}^{M}\tilde{G}_{k_j}(z)=\tilde{G}_{k_m}(z)+\sum_{j=1}^{m-1}\tilde{G}_{k_j}(z)\le\tilde{G}_{k_m}(z)+1.$$

Then we have that

$$\sum_{j=1}^{M}\tilde{G}_{k_j}\le\sum_{j=1}^{M}\tilde{G}_{k_j}\chi_{A_j}+1,$$

which combined with (2.1) and (2.3) yields

$$\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B)\le\frac{1}{\lambda}\left(\int_X|f|\,d\mu+\sum_{j=1}^{M}\int_X|f|\chi_{A_j}\tilde{G}_{k_j}\,d\mu\right).$$

In order to bound the last sum note that

$$\sum_{j=1}^{M}\int_X f\chi_{A_j}\tilde{G}_{k_j}\,d\mu=\sum_{j=1}^{M}\sum_{B\in\tilde{\mathcal{A}}_{k_j}}\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\,\mu(D_B)$$

Given $B\in\tilde{\mathcal{A}}_{k_j}$, we say that $B\in\tilde{\mathcal{A}}^*_{k_j}$ if

$$\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\le\frac{\lambda}{2}.$$

Hence, we have

$$\frac{1}{\lambda}\sum_{j=1}^{M}\sum_{B\in\tilde{\mathcal{A}}^*_{k_j}}\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\;\mu(D_B)\leq\frac{1}{2}\sum_{B\in\tilde{\mathcal{A}}}\mu(D_B)$$

and this term can be absorbed in the left hand side of (2.2).

We now consider those balls that do not belong to any of the classes $\tilde{\mathcal{A}}^*_k$. We claim that if $z\in B^0$, then $\tilde{B}\subset B(z,R_B)\subset B^*$. The n-micro-doubling condition would imply then that $\mu(B)\geq\mu(B^*)/K_0\geq\mu(B(z,R_B))/K_0$ and consequently if $B\in\tilde{\mathcal{A}}_{k_j}\setminus\tilde{\mathcal{A}}^*_{k_j}$

$$\begin{aligned}\frac{\lambda}{2} &< \frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\leq\frac{K_0}{\mu(B(z,R_B))}\int_{B(z,R_B)}|f|\chi_{A_j}\,d\mu\\ &\leq K_0M_{k_j}(f\chi_{A_j})(z).\end{aligned}$$

Thus, for such B,

$$B^0\subset\left\{x\in X: M_{k_j}(f\chi_{A_j})(x)>\frac{\lambda}{2K_0}\right\}. \tag{2.4}$$

Therefore, using that for $B\in\mathcal{A}$ one has

$$\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\leq\frac{1}{\mu(B)}\int_{B}|f|\chi_{A_j}\,d\mu\leq 2\lambda,$$

we conclude that

$$\begin{aligned}&\frac{1}{\lambda}\sum_{j=1}^{M}\sum_{B\in\tilde{\mathcal{A}}_{k_j}\setminus\tilde{\mathcal{A}}^*_{k_j}}\frac{1}{\mu(B)}\int_{\tilde{B}}|f|\chi_{A_j}\,d\mu\;\mu(D_B)\\ &\qquad\leq 2\sum_{j=1}^{M}\sum_{B\in\tilde{\mathcal{A}}_{k_j}\setminus\tilde{\mathcal{A}}^*_{k_j}}\mu(D_B)\\ &\qquad\leq 2\sum_{j=1}^{M}\mu\left(\left\{x\in X: M_{k_j}(f\chi_{A_j})(x)>\frac{\lambda}{2K_0}\right\}\right).\end{aligned}$$

This finishes the proof, provided we justify the last claim.

In order to do so, suppose that $y\in\tilde{B}$, then

$$|y-z|\leq|y-z_B|+|z_B-z|<R_{\tilde{B}}+(R_B-R_{\tilde{B}})=R_B,$$

which means that $y\in B(z,R_B)$. Assume now that $y\in B(z,R_B)$

$$|y-z_B|\leq|y-z|+|z-z_B|<R_B+R_B/n=(1+1/n)R_B,$$

hence $y\in B^*$. The claim is proved. □

3. UNIFORM BOUNDS FOR MAXIMAL FUNCTIONS ASSOCIATED WITH ROTATION INVARIANT MEASURES IN EUCLIDEAN SPACES

In the sequel we will consider X to be the Euclidean space $\mathbb{R}^n$, equipped with the Euclidean distance and a rotation invariant measure with a radial density, so that it can be defined in all dimensions. The goal of this section is to show that, in this setting, uniformly weakly doubling measures have associated maximal operators satisfying dimension free bounds on the spaces L^p.

The notation that we will use is the following. By μ we will always denote a rotation invariant measure with a radial density w. That is $d\mu(x) = w(x)\,dx$ and $w(x) = w_0(|x|)$ with $w_0 : [0,\infty) \to [0,\infty)$. Properly speaking, w and $d\mu$ are objects that change with the dimension. In particular, their integrability properties may be different from one Euclidean space to another. By a slight abuse of notation, we keep, however the same symbols to denote them in $\mathbb{R}^n$ for all n.

As usual, for a measurable set $E \subset \mathbb{R}^n$, $|E|$ will denote its Lebesgue measure, $d\sigma_{n-1}$ will be the measure on $\mathbb{S}^{n-1}$ induced by Lebesgue measure on $\mathbb{R}^n$ and $\omega_{n-1} = \sigma_{n-1}(\mathbb{S}^{n-1})$. Also, for each $x \in \mathbb{R}^n$ and $R > 0$, we will denote by $B(x,R)$ the Euclidean ball centered at x with radius R. For a ball whose center is the origin we will use the notation B_R.

3.1. Weakly-doubling measures in Euclidean spaces and uniform bounds.

As a first and simple example that uniformity in the weak doubling condition implies uniform bounds we present

Theorem 3.1. *Let μ be a rotation invariant measure over $\mathbb{R}^n$ that is absolutely continuous with respect to Lebesgue measure. If μ is weakly doubling with constant K_1 and f is a radial function we have*

$$\mu(\{x \in \mathbb{R}^n : M_\mu f(x) > \lambda\}) \le \frac{2(K_1+1)}{\lambda}\|f\|_{L^1(\mathbb{R}^n,\mu)}.$$

Observe that if K_1 is uniformly bounded in n, the above inequality provides a dimension-free bound for M_μ. This is a generalization of a result presented by M.T. Menárguez and the second author in [25]. There, it was shown that for all radial f over $\mathbb{R}^n$ and $\lambda > 0$ one has

$$|\{x \in \mathbb{R}^n : Mf(x) > \lambda\}| \le \frac{4}{\lambda}\|f\|_{L^1(\mathbb{R}^n,dx)}. \tag{3.1}$$

A previous generalization was due to A. Infante, who pointed out (see [20], [21]) that the same is true for M_μ, if μ is an 'increasing' measure. By this we mean that $\mu(B(x,R)) \ge \mu(B(y,R))$ whenever $y = ax$ with $a \ge 1$. In [4] there is an independent approach with results similar to our Theorems 3.1 and 4.3. These two results already appeared in [14].

The main result of the section asserts that the uniform weakly-doubling property of a measure μ ensures uniform bounds in the dimension not only for radial functions but, in fact, over the whole class of functions $L^p(\mathbb{R}^n, d\mu)$, if $p > 1$.

Theorem 3.2. *If there is a positive integer N so that μ is uniformly weakly-doubling in $\mathbb{R}^n$ for $n \ge N$, then for each $p > 1$ there exists C_p only depending on μ and p so that*

$$\|M_\mu f\|_{L^p(\mathbb{R}^n,\mu)} \le C_p\|f\|_{L^p(\mathbb{R}^n,\mu)},$$

for all $f \in L^p(\mathbb{R}^n, \mu)$.

In order to prove this result we first obtain the following characterization of radial measures that are uniformly weakly-doubling.

Theorem 3.3. *There exists $N \in \mathbb{N}$ so that $d\mu = w(x)dx$, with $w(x) = w_0(|x|)$, is uniformly weakly-doubling in $\mathbb{R}^n$ for each $n \geq N$ if and only if w is essentially constant on dyadic annuli, i.e. there exists $\beta \geq 1$ so that for all $R > 0$ one has*

$$\operatorname*{ess\,sup}_{R \leq |x| \leq 2R} w(x) \leq \beta \operatorname*{ess\,inf}_{R \leq |x| \leq 2R} w(x). \tag{3.2}$$

The proof of this employs a new method of differentiation through dimensions that is presented in Section 3.2. With this characterization at hand, instead of proving Theorem 3.2 we will prove the equivalent:

Theorem 3.4. *With the above notation, let w be essentially constant over dyadic annuli with constant β. Then, there exists N so that μ is locally finite in $\mathbb{R}^n$ for $n \geq N$ and, for each $p > 1$, there exists a constant C_p only depending on p and β so that for $n \geq N$ one has*

$$\|M_\mu f\|_{L^p(\mathbb{R}^n,\mu)} \leq C_p \|f\|_{L^p(\mathbb{R}^n,\mu)},$$

for all $f \in L^p(\mathbb{R}^n, \mu)$.

The densities $|x|^\alpha$, $\alpha \in \mathbb{R}$, provide examples of measures with this property. It is easy to see that in each case they are essentially constant over dyadic annuli with constant $2^{|\alpha|}$.

At this point, let us make the simple observation that if w is essentially constant over dyadic annuli, then w can be compared pointwise with a continuous function in $\mathbb{R}^n \setminus \{0\}$. In the sequel, we will assume, as we may, that under this hypothesis, w is indeed continuous with the possible exception at $x = 0$.

Theorem 3.4 will be obtained as a consequence of the following results.

Lemma 3.5. *Let μ be locally finite in $\mathbb{R}^n$ and assume that its density w is essentially constant over dyadic intervals with constant β. Then we have the following pointwise inequality*

$$M_\mu f(x) \leq C \left(Mf(x) + \frac{1}{w(x)} M(fw)(x) + \mathcal{H}_\mu f(x) \right),$$

where C depends only on β, M is the usual Hardy-Littlewood maximal operator and $\mathcal{H}_\mu$ is defined as

$$\mathcal{H}_\mu f(x) = \sup_{R \geq |x|} \frac{1}{\mu(B_R)} \int_{B_R} |f(y)| \, d\mu(y).$$

This transforms the problem of finding $L^p(\mu)$ bounds for M_μ to the one of finding them for $\mathcal{H}_\mu$ and for M.

Theorem 3.6. *In the same hypotheses of Lemma 3.5, assume that in addition we have the weighted inequalities*

$$\begin{array}{rcl} \|Mf\|_{L^p(w)} & \leq & W_1 \|f\|_{L^p(w)}, \\ \|Mf\|_{L^p(w^{1-p})} & \leq & W_2 \|f\|_{L^p(w^{1-p})}. \end{array} \tag{3.3}$$

Then one has

$$\|M_\mu f\|_{L^p(\mu)} \leq C \|f\|_{L^p(\mu)},$$

where the constant C only depends on β, W_1 and W_2.

Note that the required weighted inequalities are equivalent to $w \in A_p \cap A_{p'} = A_{\min\{p,p'\}}$, where $A_p = A_p(\mathbb{R}^n)$ denotes the usual Muckenhoupt class of weights. The existence of W_1 and W_2 in (3.3) is guaranteed due to the following

Lemma 3.7. *Let w be essentially constant over dyadic intervals with constant β. Then for each $p > 1$ there exist $N \in \mathbb{N}$ depending only on p and β so that $w \in A_p(\mathbb{R}^n)$ for all $n \geq N$.*

The constants W_1 and W_2 can be taken, in fact, independent of the dimension, as the following result by J. Duoandikoetxea and L. Vega, appeared in [16], shows.

Theorem 3.8. *Let w_0 be a nonnegative function on $(0, \infty)$, so that $w = w_0(|\cdot|) \in A_p(\mathbb{R}^N)$. Then for all $n \geq N$ one has $w \in A_p(\mathbb{R}^n)$. Moreover,*

$$\|Mf\|_{L^p(\mathbb{R}^n,w)} \leq C\|f\|_{L^p(\mathbb{R}^n,w)},$$

with a constant C that might depend on p and w but not on n.

To complete the proof of Theorem 3.4, we have to show that if w is essentially constant over dyadic annuli, then there exists $N \in \mathbb{N}$ so that w is locally integrable in $\mathbb{R}^n$ for $n > N$. This is immediate once we have the following auxiliary lemma, which will be used in different proofs throughout the paper.

Lemma 3.9. *Let w be essentially constant over dyadic intervals with constant β. Then there exists $N \in \mathbb{N}$ only depending on β, so that for $n \geq N$ one has in $\mathbb{R}^n$ the control*

$$\mu(B_R) = \int_{B_R} w(x)\,dx \leq 2\beta\, w_0(R)|B_R|. \tag{3.4}$$

In the remaining part of the section we present the proofs of Theorem 3.6, Lemmas 3.5, 3.7 and 3.9 and Theorem 3.1 in this given order.

We start showing how Lemma 3.5 implies Theorem 3.6.

Proof of Theorem 3.6. By Lemma 3.5 one has

$$\|M_\mu f\|_{L^p(\mu)} \leq C\left(\|Mf\|_{L^p(\mu)} + \left\|\frac{M(fw)}{w}\right\|_{L^p(\mu)} + \|\mathcal{H}_\mu f\|_{L^p(\mu)}\right),$$

where the constant C depends only on β.

By assumption we have

$$\|Mf\|_{L^p(\mu)} \leq W_1\|f\|_{L^p(\mu)}.$$

We also would like to have

$$\left\|\frac{1}{w}M(fw)\right\|_{L^p(\mu)} \leq C\|f\|_{L^p(\mu)}.$$

Taking $g = fw$ this is equivalent to

$$\|Mg\|_{L^p(w^{1-p})} \leq C\|g\|_{L^p(w^{1-p})},$$

which we know true by assumption with $C = W_2$.

For $\mathcal{H}_\mu$ we use the standard argument for Hardy type operators. It is obvious that $\mathcal{H}_\mu$ is bounded on $L^\infty(\mu)$ with constant 1. We will show that it is also weakly bounded on $L^1(\mu)$ with operator norm 1. Then by real interpolation it is bounded on $L^p(\mu)$ with operator norm controlled by an absolute constant.

To see the weak type inequality take $\lambda > 0$ and consider $E_\lambda = \{x \in \mathbb{R}^n : \mathcal{H}_\mu f(x) \geq \lambda\}$. If $x \in E_\lambda$ there exists $R_x > |x|$ so that

$$\frac{1}{\mu(B_{R_x})} \int_{B_{R_x}} |f(y)|\, d\mu(y) \geq \lambda.$$

Note that then $B_{R_x} \subset E_\lambda$, and that

$$E_\lambda = \bigcup_{x \in E_\lambda} B_{R_x}.$$

Then $E_\lambda = B_R$ for certain $R > 0$ and monotonicity gives

$$\begin{aligned}
\mu(E_\lambda) &= \mu(B_R) = \sup_{x \in E_\lambda} \mu(B_{R_x}) \leq \sup_{x \in E_\lambda} \frac{1}{\lambda} \int_{B_{R_x}} |f(y)|\, d\mu(y) \\
&\leq \frac{1}{\lambda} \int_{\mathbb{R}^n} |f(y)|\, d\mu(y).
\end{aligned}$$

□

Proof of Lemma 3.5. We will bound the mean value over an arbitrary ball $B(x,R)$. Fixing x and R, we consider different cases.

If $|x| \geq 2R$ and $y \in B(x,R)$ then

$$\frac{1}{2}|x| \leq |x| - R \leq |y| \leq |x| + R \leq \frac{3}{2}|x|.$$

Since w is essentially constant over dyadic annuli, we have $\beta^{-1}w(x) \leq w(y) \leq \beta w(x)$. Hence

$$\begin{aligned}
\frac{1}{\mu(B(x,R))} \int_{B(x,R)} |f(y)|\, w(y)\, dy &\leq \frac{\beta^2}{w(x)\, |B(x,R)|} \int_{B(x,R)} |f(y)|\, w(x)\, dy \\
&\leq \beta^2 M f(x).
\end{aligned}$$

In the case that $R/2 \leq |x| \leq 2R$ one has that if $y \in B(x,R) \setminus B_{R/2}$, then $|x|/4 \leq |y| \leq 3|x|$, which implies $\beta^{-2}w(|x|) \leq w(y) \leq \beta^2 w(|x|)$. Hence,

$$\begin{aligned}
\mu(B(x,R)) &\geq \mu(B(x,R) \setminus B_{R/2}) \geq \frac{1}{\beta^2}\, w(x)\, |B(x,R) \setminus B_{R/2}| \\
&\geq \frac{1}{2\beta^2}\, w(x)\, |B(x,R)|.
\end{aligned}$$

Therefore we have

$$\begin{aligned}
\frac{1}{\mu(B(x,R))} \int_{B(x,R)} |f(y)|\, w(y)\, dy &\leq \frac{2\beta^2}{w(x)|B(x,R)|} \int_{B(x,R)} |f(y)|\, w(y)\, dy \\
&\leq \frac{2\beta^2}{w(x)} M(fw)(x).
\end{aligned}$$

Last, we consider that $|x| \leq R/2$. We split the ball $B(x,R)$ into two disjoint pieces $B_{R/2}$ and $B(x,R) \setminus B_{R/2}$ and integrate over them separately. For the first one

$$\frac{1}{\mu(B(x,R))} \int_{B_{R/2}} |f(y)|\, d\mu(y) \leq \frac{1}{\mu(B_{R/2})} \int_{B_{R/2}} |f(y)|\, d\mu(y) \leq \mathcal{H}_\mu f(x).$$

For the second one note that if $y \in B(x,R) \setminus B_{R/2}$ then $R/2 \leq |y| \leq 3R$ and then $\beta^{-2} w_0(R) \leq w(y) \leq \beta^2 w_0(R)$. This implies that

$$\begin{aligned} \mu(B(x,R)) &\geq \mu(B(x,R) \setminus B_{R/2}) \geq \frac{1}{\beta^2}\, w_0(R)\, |B(x,R) \setminus B_{R/2}| \\ &\geq \frac{1}{2\beta^2}\, w_0(R)\, |B(x,R)|, \end{aligned}$$

and then we get

$$\begin{aligned} \frac{1}{\mu(B(x,R))} & \int_{B(x,R)\setminus B_{R/2}} |f(y)|\, d\mu(y) \\ &\leq \frac{2\beta^4}{w_0(R)\, |B(x,R)|} \int_{B(x,R)\setminus B_{R/2}} |f(y)|\, w_0(R)\, dy \\ &\leq 2\beta^4 M f(x). \end{aligned}$$

□

Remark. Both, Theorem 3.6 and Lemma 3.5 do not require μ to be radial. For the applications, however, this requirement is the most natural in order to define μ and M_μ simultaneously in all dimensions and to study whether or not there are uniform bounds as $n \longrightarrow \infty$.

Proof of Lemma 3.7. We have to prove that $w \in A_p(\mathbb{R}^N)$ for some $N \in \mathbb{N}$. That is, there exists a constant $C > 0$ so that for all $x \in \mathbb{R}^N$ and $R > 0$ one has

$$⨍_{B(x,R)} w(y)\, dy \left(⨍_{B(x,R)} w(y)^{1/(1-p)}\, dy \right)^{p-1} \leq C.$$

Observe that $w^{1/(1-p)}$ is also constant over dyadic annuli. This is easy because

$$\begin{aligned} \sup_{R \leq |x| \leq 2R} w(x)^{1/(1-p)} &= \left(\inf_{R \leq |x| \leq 2R} w(x) \right)^{1/(1-p)} \\ &\leq \left(\beta^{-1} \sup_{R \leq |x| \leq 2R} w(x) \right)^{1/(1-p)} \\ &= \beta^{1/(p-1)} \inf_{R \leq |x| \leq 2R} w(x)^{1/(1-p)}. \end{aligned}$$

Then we can choose N so that (3.4) holds for w and $w^{1/(1-p)}$ in $\mathbb{R}^N$. If $|x| \le 2R$, one has $B(x,R) \subset B_{3R}$ and then

$$\begin{aligned} &\fint_{B(x,R)} w(x)\,dx \, \left(\fint_{B(x,R)} w(x)^{1/(1-p)}\,dx\right)^{p-1} \\ &\qquad \le \; 3^{pn} \fint_{B_{3R}} w(x)\,dx \left(\fint_{B_{3R}} w(x)^{1/(1-p)}\,dx\right)^{p-1} \\ &\qquad \le \; 3^{pn} 2\beta\, w_0(3R) \left(2\beta^{1/(p-1)}\, w_0(3R)^{1/(1-p)}\right)^{p-1} \\ &\qquad \le \; (2\,3^n)^p \beta^2. \end{aligned}$$

Assume conversely that $|x| > 2R$. If $y \in B(x,R)$, then $|x|/2 \le |y| \le 3|x|/2$ and consequently

$$\beta^{-1} w(x) \le w(y) \le \beta w(x).$$

Hence,

$$\begin{aligned} &\fint_{B(x,R)} w(y)\,dy \, \left(\fint_{B(x,R)} w(y)^{1/(1-p)}\,dy\right)^{p-1} \\ &\qquad \le \; \beta w(x) \left((\beta^{-1} w(x)\,)^{1/(1-p)}\right)^{p-1} \le \beta^2. \end{aligned}$$

□

Proof of Lemma 3.9. Assuming that we are in $\mathbb{R}^N$ with $2^N > \beta$ we have

$$\begin{aligned} \mu(B_R) &= \omega_{N-1} \int_0^R w_0(t) t^{N-1}\,dt = \omega_{N-1} \sum_{j=0}^{\infty} \int_{2^{-j-1}R}^{2^{-j}R} w(t) t^{N-1}\,dt \\ &\le \omega_{N-1} \sum_{j=0}^{\infty} \int_{2^{-j-1}R}^{2^{-j}R} \beta^{j+1} w_0(R)\, t^{N-1}\,dt \\ &\le \beta\, w_0(R) \frac{\omega_{N-1} R^N}{N} \sum_{j=0}^{\infty} \left(\frac{\beta}{2^N}\right)^j = \frac{\beta}{1-\beta/2^N}\, w_0(R)\, |B_R|. \end{aligned}$$

Taking N big enough one has that $1/(1-\beta/2^N) \le 2$, as stated. □

For the proof of Theorem 3.1, we will follow [25]. First, associated with a weight v on $\mathbb{R}$ we define the non-centered maximal function

$$\tilde{M}_v F(x) = \sup_{a \le x \le b} \frac{1}{v([a,b])} \int_{[a,b]} |F(t)| v(t)\,dt, \tag{3.5}$$

for each F locally integrable with respect to v. Then, using a simple covering argument for intervals in $\mathbb{R}$ one has the estimate

$$v(\{r \ge 0 \,:\, \tilde{M}_v F(r) > \lambda\}) \le \frac{2}{\lambda} \int_{\mathbb{R}} |F(r)| v(r)\,dr, \tag{3.6}$$

(see [26], [18] or [23]).

We make now the following definition. Given a measurable set $E \in \mathbb{R}^n$ we define its projection onto the sphere $\mathbb{S}^{n-1}$ by

$$\Sigma_E = \{\theta \in \mathbb{S}^{n-1} \;:\; r\theta \in E \text{ for some } r > 0\}.$$

The following geometrical result can be found in [25].

Lemma 3.10. *For each ball* $B(x,R)\in\mathbb{R}^n$ *there exists a set* D *such that*

(a) $B(x,R)\subset D$,
(b) $\Sigma_D=\Sigma_{B(x,R)}$,
(c) for each $\theta\in\Sigma_D$ *there exist* $0\le a_\theta\le b_\theta$ *such that* $r\theta\in\bar{D}$ *if and only if* $a_\theta\le r\le b_\theta$,
(d) $|x|\theta\in\bar{D}$ *for each* $\theta\in\sigma_D$ *(this means* $a_\theta\le|x|\le b_\theta$*),*
(e) D *is contained in the union of* $B(x,R)$ *and another ball* $B(z,R)$ *with* $z\in B(x,R)$.

Using this, we prove Theorem 3.1.

Proof of Theorem 3.1. By conditions *(a)* and *(e)* and the hypothesis that μ is weakly doubling, we have

$$\begin{aligned}\frac{1}{\mu(B(x,R))}\int_{B(x,R)}|f(y)|\,d\mu(y) &\le \frac{\mu(D)}{\mu(B(x,R))}\frac{1}{\mu(D)}\int_D|f(y)|\,d\mu(y)\\ &\le \frac{1+K_1}{\mu(D)}\int_D|f(y)|\,d\mu(y).\end{aligned}$$

Assume that $f(x)=f_0(|x|)$ and set $v(t)=w_0(t)t^{n-1}$. Now we integrate along each ray coming from the origin and use conditions *(b)–(d)*

$$\begin{aligned}\int_D|f(y)|\,d\mu(y) &= \int_{\Sigma_D}\int_{a_\theta}^{b_\theta}|f_0(t)|v(t)\,dt\,d\sigma_n(\theta)\\ &= \int_{\Sigma_D}\frac{v([a_\theta,b_\theta])}{v([a_\theta,b_\theta])}\int_{a_\theta}^{b_\theta}|f_0(t)|v(t)\,dt\,d\sigma_n(\theta)\\ &\le \int_{\Sigma_D}v([a_\theta,b_\theta])\,\tilde{M}_vf_0(|x|)\,d\sigma_n(\theta).\end{aligned}$$

Note that

$$\int_{\Sigma_D}v([a_\theta,b_\theta])\,d\sigma_n(\theta)=\int_{\Sigma_D}\int_{a_\theta}^{b_\theta}v(t)\,dt\,d\sigma_n(\theta)=\mu(D),$$

Hence we have proved that

$$\frac{1}{\mu(B(x,R))}\int_{B(x,R)}|f(y)|\,d\mu(y)\le(1+K_1)\,\tilde{M}_vf_0(|x|),$$

and, since R is arbitrary, $M_\mu f(x)\le(1+K_1)\tilde{M}_vf_0(|x|)$. Integrating in polar coordinates we have

$$\mu\left(\{x\in\mathbb{R}^n\ :\ Mf(x)>\lambda\}\right)\le\omega_{n-1}\,v\left(\left\{r\ge 0\ :\ \tilde{M}_vf_0(r)>\frac{\lambda}{1+K_1}\right\}\right).$$

By (3.6) the latter term is bounded by

$$\frac{2(1+K_1)}{\lambda}\,\omega_{n-1}\int_0^\infty|f_0(t)|\,v(t)\,dt=\frac{2(K_1+1)}{\lambda}\int_{\mathbb{R}^n}|f(x)|\,d\mu(x).$$

□

3.2. **Differentiation through dimensions.** In this section we introduce a new technique that will be useful in settling several questions mentioned in this work, among them, the proof of Theorem 3.3. It starts with the following observation: take, in each Euclidean space $\mathbb{R}^n$, a ball B^n of a fixed radius R and with center at a fixed distance s from the origin. If $w_0 \in L^1_{\text{loc}}([0,\infty), t^{N-1}dt)$, for some $N \geq 1$, and m_n denotes Lebesgue measure in $\mathbb{R}^n$, then the limits

$$\lim_{n\to\infty} ⨍_{B^n} w_0(|x|)\, dm_n(x)$$

exist a.e. depending on R and s. Observe that our integrability condition on w_0 ensures that the function $w(x) = w_0(|x|)$ is locally integrable in each $\mathbb{R}^n$ whenever $n \geq N$. Let us remark at this point that for balls B^n of the same radius one always has $\lim_{n\to\infty} m_n(B^n) = 0$. This is not however the reason for this phenomenon that we will refer to as "differentiation through dimensions". The precise statement of this type of differentiation is contained in the following result.

Lemma 3.11. *Take $w_0 \in L^1_{\text{loc}}([0,\infty), t^{N-1}\, dt)$ for some $N \geq 1$. Then, for almost every $T > 0$ and for all $s \geq 0$ and $R > 0$ so that $s^2 + R^2 = T^2$, if we take points $z^n \in \mathbb{R}^n$ with $|z^n| = s$ and we denote $B(z^n, R) = \{y \in \mathbb{R}^n : |z^n - y| < R\}$, the following holds*

$$\lim_{n\to\infty} ⨍_{B(z^n,R)} w_0(|x|)\, dm_n(x) = w_0(T).$$

Proof. The idea is to exploit the fact that in high dimensions the measure of a ball concentrates around 'maximal circles'. We will assume that w_0 is positive.

Fix $T > 0$, and take positive s, $z^n \in \mathbb{R}^n$ with $|z^n| = s$ and R with $s^2 + R^2 = T^2$. Observe that

$$⨍_{B(z^n,R)} w_0(|x|)\, dm_n(x) = \frac{n}{\omega_{n-1}R^n} \int_{(s-R)_+}^{s+R} w_0(t) A_n(t) t^{n-1}\, dt.$$

where $A_n(t) = A_n^{(s,R)}(t) = \left|\{\theta \in \mathbb{S}^{n-1} : t\theta \in B(z^n, R)\}\right|_{n-1}$. Define

$$\varphi_n(t) = \varphi_n^{(s,R)}(t) = \frac{n}{\omega_{n-1}R^n} A_n(t) t^{n-1} \chi_{[(s-R)_+, s+R]}(t).$$

With this notation we have

$$⨍_{B(z^n,R)} w_0(|x|)\, dm_n(x) = \int_{-\infty}^{\infty} w_0(t) \varphi_n^{(s,R)}(t)\, dt =: \Phi_n^{(s,R)} w_0.$$

For a continuous w_0 the proof follows if we show that $\varphi_n \overset{n\to\infty}{\longrightarrow} \delta_T$ in the sense of distributions. For this it is enough to check that φ_n is an approximation of the identity at the point $t = T$, that is

i) $\varphi_n \geq 0$ for all $n \in \mathbb{N}$,
ii) $\int_{-\infty}^{\infty} \varphi_n(t)\, dt = 1$ for all $n \in \mathbb{N}$,
iii) $\forall \varepsilon > 0$ one has $I_\varepsilon(n) = \int_{|t-T|>\varepsilon} \varphi_n(t)\, dt \overset{n\to\infty}{\longrightarrow} 0$.

Note that i) is trivial and ii) is immediate from the observation that

$$\omega_{n-1} \int_{(s-R)_+}^{s+R} A_n(t) t^{n-1}\, dt = |B(z^n, R)|_n.$$

To see iii) observe that $I_\varepsilon(n) = I^1_\varepsilon(n) + I^2_\varepsilon(n)$ where

$$\begin{aligned} I^1_\varepsilon(n) &= \int_{-\infty}^{T-\varepsilon} \varphi_n(t)\, dt = \frac{|B(z^n,R) \cap B_{T-\varepsilon}|_n}{|B(z^n,R)|_n} \leq \left(\frac{R_1}{R}\right)^n \overset{n\to\infty}{\longrightarrow} 0, \\ I^2_\varepsilon(n) &= \int_{T+\varepsilon}^{\infty} \varphi_n(t)\, dt = \frac{|B(z^n,R) \setminus B_{T+\varepsilon}|_n}{|B(z^n,R)|_n} \leq \left(\frac{R_2}{R}\right)^n \overset{n\to\infty}{\longrightarrow} 0, \end{aligned}$$

and R_1 and R_2 are the radii of the minimal balls that contain $B(z^n,R)\cap B_{T-\varepsilon}$ and $B(z^n,R)\setminus B_{T+\varepsilon}$ respectively. It is obvious that $R_1,R_2<R$. This proves Lemma 3.11, $\forall T>0$, if w_0 is continuous in $[0,\infty)$.

For a general w_0 we have to show that, if $P_T=\{(s,R)\,:\,s\geq 0,\ R>0,\ s^2+R^2=T^2\}$, the set $E_0(w_0)$ defined as

$$\left\{T\in[\varepsilon,T_0]\,:\,\exists(s,R)\in P_T\,:\,\limsup_{n\to\infty}\Phi_n^{(s,R)}w_0-\liminf_{n\to\infty}\Phi_n^{(s,R)}w_0>0\right\},$$

has measure 0 for every ε,T_0 with $0<\varepsilon<T_0$.

To that end, we fix $T\in[\varepsilon,T_0]$. For $t\in[(s-R)_+,s+R]$ we denote by $y(t)$ the diameter of the set $\partial B_t\cap B(z^n,R)$. Observe that $y(t)\leq R$ and that y increases with t up to the point $t=T$, where it attains its maximum, and then decreases in $(T,s+R)$. Also, for each $t\in[|s-R|,s+R]$ we call $\alpha(t)$ the angle between the segment connecting the origin with z^n and the one joining the origin with any point in $\partial B_t\cap\partial B(z^n,R)$. Clearly $y(t)=t\sin\alpha(t)$.

Observe first that the function $\varphi_n^{(s,R)}$ is decreasing in the interval $[T,s+R]$. To see this note that

$$\frac{A_n(t)t^{n-1}}{\omega_{n-2}}=t^{n-1}\int_0^{\alpha(t)}(\sin\beta)^{n-2}\,d\beta=\int_0^{y(t)}u^{n-2}\,\frac{du}{\sqrt{1-(u/t)^2}},\tag{3.7}$$

and that, both, the integrand and $y(t)$ decrease with t in this interval. Hence,

$$\int_T^{s+R}\varphi_n^{(s,R)}(t)w_0(t)\,dt\leq M_{\rightarrow}\overline{w}_0(T),$$

where $M_{\rightarrow}$ denotes the one-sided maximal operator

$$M_{\rightarrow}f(t)=\sup_{h>0}\frac{1}{h}\int_t^{t+h}|f(s)|\,ds,$$

and $\overline{w}_0$ is the restriction of w_0 to the interval $[\varepsilon,2T_0]$.

Let us assume that $s>0$. If $\frac{\pi}{2}\leq\alpha(t)\leq\pi$, we will use that

$$t^{1-N}\,\varphi_n^{(s,R)}(t)\leq\frac{nt^{n-N}}{R^n}.\tag{3.8}$$

This case may happen only if $R>s$ and, then, $t\leq\sqrt{R^2-s^2}<R$. For $\alpha(t)<\frac{\pi}{2}$ we have the estimate

$$t^{1-N}\,\varphi_n^{(s,R)}(t)\leq\frac{nt^{n-N}}{\omega_{n-1}R^n}\,\omega_{n-2}\,\alpha(t)\,\sin^{n-2}\alpha(t)\leq\frac{n\omega_{n-2}}{\omega_{n-1}R^n}\,\frac{\pi}{2}\,y(t)^{n-N}.\tag{3.9}$$

Combining (3.8) and (3.9) we see that for all $t_0<T$ we have

$$\lim_{n\to\infty}\left[\sup_{t<t_0}t^{1-N}\,\varphi_n^{(s,R)}(t)\right]=0.$$

As a consequence, if $t_0<T$

$$\begin{aligned}\limsup_{n\to\infty}\int_0^{t_0}\varphi_n^{(s,R)}(t)\,w_0(t)\,dt&\leq\lim_{n\to\infty}\left[\sup_{t<t_0}t^{1-N}\,\varphi_n^{(s,R)}(t)\right]\int_0^{t_0}w_0(t)t^{N-1}\,dt\\&=0.\end{aligned}$$

Still for $\alpha(t)<\frac{\pi}{2}$, we make the observation that

$$A_n(t)=\omega_{n-2}\int_0^{\alpha(t)}(\sin\beta)^{n-2}\,\cos\beta\,\cos^{-1}\beta\,d\beta,$$

and so, $\frac{\omega_{n-2}}{n-1}\,\sin^{n-1}\alpha(t)\le A_n(t)\le \frac{\omega_{n-2}}{n-1}\,\sin^{n-1}\alpha(t)\,\cos^{-1}\alpha(t)$. Thus, if we call

$$\tilde{\varphi}_n^{(s,R)}(t)=\frac{n}{n-1}\frac{\omega_{n-2}}{\omega_{n-1}}\frac{y(t)^{n-1}}{R^n},$$

then we have the double estimate

$$\tilde{\varphi}_n^{(s,R)}(t)\le \varphi_n^{(s,R)}(t)\le \tilde{\varphi}_n^{(s,R)}(t)\cdot\cos^{-1}\alpha(t).$$

It is important to point out that $\tilde{\varphi}_n^{(s,R)}(t)$ is increasing in $[0,T]$ and that

$$\int_{[0,T]}\tilde{\varphi}_n^{(s,R)}(t)\,dt\le\int_{[0,T]}\varphi_n^{(s,R)}(t)\,dt\le 1.$$

The final observation is that there exists $t_1=t_1(s,R)<T$, so that $\alpha(t)$ decreases in $[t_1,T]$. Hence, for $t_0\in[\max\{t_1,\,\varepsilon\},\,T)$ we have

$$\begin{aligned}\int_{t_0}^{T}\varphi_n^{(s,R)}(t)w_0(t)\,dt &\le \int_{t_0}^{T}\tilde{\varphi}_n^{(s,R)}(t)w_0(t)\,\cos^{-1}\alpha(t)\,dt\\ &\le \cos^{-1}\alpha(t_0)M_{\leftarrow}\overline{w}_0(T),\end{aligned}$$

where $M_{\leftarrow}$ is the maximal operator

$$M_{\leftarrow}f(t)=\sup_{h>0}\frac{1}{h}\int_{t-h}^{t}|f(s)|\,ds.$$

For this t_0,

$$\begin{aligned}\limsup_{n\to\infty}\Phi_n^{(s,R)}w_0 &= \limsup_{n\to\infty}\left(\int_0^{t_0}\varphi_n^{(s,R)}(t)w_0(t)\,dt+\int_{t_0}^{T}\varphi_n^{(s,R)}(t)w_0(t)\,dt\right)\\ &\le \cos^{-1}\alpha(t_0)M_{\leftarrow}\overline{w}_0(T).\end{aligned}$$

Taking the limit as $t_0\to T^-$, we have

$$\begin{aligned}\limsup_{n\to\infty}\Phi_n^{(s,R)}w_0 &\le \cos^{-1}\alpha(T)M_{\leftarrow}\overline{w}_0(T)=\frac{T}{s}\,M_{\leftarrow}\overline{w}_0(T)\\ &\le \left(1+\frac{R}{s}\right)\,M_{\leftarrow}\overline{w}_0(T).\end{aligned}$$

Therefore, if $R/s\le k$ for $k\in\mathbb{N}$ we have

$$\limsup_{n\to\infty}\Phi_n^{(s,R)}w_0\le M_{\rightarrow}\overline{w}_0(T)+(1+k)M_{\leftarrow}\overline{w}_0(T)\le 3k\tilde{M}\overline{w}_0(T),$$

with $\tilde{M}$ denoting, as in (3.5), the non centered one dimensional maximal operator

$$\tilde{M}f(t)=\sup_{a\le t\le b}\frac{1}{b-a}\int_a^b|f(s)|\,ds.$$

Observe that if $s=0$, then $\varphi_n^{(0,T)}(t)$ is an increasing function in the interval $[0,T]$. Arguing as before to remove the integration on $[0,\varepsilon]$, we have

$$\limsup_{n\to\infty}\Phi_n^{(0,T)}w_0\le M_{\leftarrow}\overline{w}_0(T)\le\tilde{M}\overline{w}_0(T).$$

To finish the proof, for $T>0$ and $k\in\mathbb{N}$ we denote $P_T^k=\{(s,R)\,:\,s,R>0,\ R/s\le k,\ s^2+R^2=T^2\}\cup\{(0,T)\}$ and we define for $\lambda>0$ the set $E_\lambda^k(w_0)$ as

$$\left\{T\in[\varepsilon,\,T_0]\,:\,\exists(s,R)\in P_T^k\,:\,\limsup_{n\to\infty}\Phi_n^{(s,R)}w_0-\liminf_{n\to\infty}\Phi_n^{(s,R)}w_0>\lambda\right\}.$$

Note that if g is continuous, the previous considerations and (3.6) give

$$\begin{aligned}\left|E_\lambda^k(w_0)\right| &= \left|E_\lambda^k(w_0-g)\right| \leq \left|\left\{T>0\,:\,6k\tilde{M}(\overline{w}_0-\overline{g})(T)>\lambda\right\}\right|\\ &\leq \frac{12k}{\lambda}\|\overline{w}_0-\overline{g}\|_{L^1},\end{aligned}$$

This last term can be made as small as needed with an appropriate choice of g. Therefore $\left|E_\lambda^k(w_0)\right| = 0$. As a consequence, $\left|E_0^k(w_0)\right| = \left|\bigcup_{\lambda>0}E_\lambda^k(w_0)\right| = 0$ and, hence, $|E_0(w_0)| = \left|\bigcup_{k\in\mathbb{N}}E_0^k(w_0)\right| = 0$, as wanted.

□

3.3. **Some applications of the differentiation through dimensions.** As a consequence of this new technique of differentiation, we prove here Theorem 3.3 and show the equivalence between the two properties of weakly doubling and strong n-micro-doubling when they hold uniformly with the dimension.

Proof of Theorem 3.3. First suppose that μ is uniformly weakly doubling with constant K_1 in each $\mathbb{R}^n$ with $n\geq N$ for some $N\in\mathbb{N}$. We have to prove that w is essentially constant over dyadic annuli.

With P_T as in the previous proof, let

$$\mathcal{T} = \left\{T>0\,:\,\lim_{n\to\infty}\frac{\mu(B(z^n,R))}{|B(z^n,R)|} = w_0(T), \quad \forall(|z^n|,R)\in P_T\right\}.$$

We know from Lemma 3.11 that $\mathcal{T}$ has full measure in $(0,\infty)$.

Now, for $R\in\mathcal{T}$ and $T\in\mathcal{T}\cap[R,2R]$ we take $z^n\in\mathbb{R}^n$ with $|z^n|^2+R^2=T^2$. Observe that $0\leq|z^n|\leq\sqrt{(2R)^2-R^2}=\sqrt{3}R$, so that $B_R\cap B(z^n,R)\neq\emptyset$. By the hypothesis on μ

$$\frac{1}{K_1}\frac{\mu(B_R)}{|B_R|}\leq\frac{\mu(B(z^n,R))}{|B_R|}\leq K_1\,\frac{\mu(B_R)}{|B_R|}.$$

Taking limits when $n\to\infty$, as we may since $R,T\in\mathcal{T}$ we get

$$\frac{1}{K_1}\,w_0(R)\leq w_0(T)\leq K_1\,w_0(R). \tag{3.10}$$

This shows that w_0 is essentially constant on dyadic intervals with constant $\beta=K_1$.

The reverse implication is straightforward and will be omitted. □

We finish this section with a proof of the following important equivalence.

Theorem 3.12. *A rotation invariant measure μ is uniformly strong n-micro-doubling in each $\mathbb{R}^n$ for $n\geq N$ if and only if μ is uniformly weakly doubling.*

Proof. In view of Theorem 3.3 we only need to prove that a measure given by a radial density that is essentially constant over dyadic annuli, is also uniformly n-micro-doubling.

Assume that w is essentially constant over dyadic annuli with constant β and take $n\geq N$ for the $N\in\mathbb{N}$ obtained in Lemma 3.9. Given $x\in\mathbb{R}^n$ and $R>0$, we will write $B^*(x,R)=B(x,(1+1/n)R)$. We first consider the case $|x|\geq 3R$, which implies that $(1+1/n)R\leq 2|x|/3$. If $y\in$

$B^*(x,R)$, then $|x|/3 \le |y| \le 5|x|/3$ which means that $\beta^{-2}\, w(x) \le w(y) \le \beta^2 w(x)$. From this we get

$$\begin{aligned} \mu\big(B^*(x,R)\big) &= \int_{B^*(x,R)} w(y)\,dy \le \beta^2\, w(x)\,|B^*(x,R)| \\ &\le e\beta^2\, w(x)\,|B(x,R)| \le e\beta^4\,\mu(B(x,R)). \end{aligned}$$

If $|x| \le 3R$ we split $B^*(x,R)$ into two disjoint pieces. For the one intersecting $B_{R/2}$ we use Lemma 3.9 as follows

$$\mu(B^*(x,R)\cap B_{R/2}) \le \mu(B_{R/2}) \le 2\beta\, w_0(R/2)\,|B_{R/2}| \le \beta^2\, w_0(R)\,|B_R|.$$

In the complementary piece w is essentially constant: if $y \in B^*(x,R)\setminus B_{R/2}$, then $R/2 \le |y| \le 5R$, which means that $\beta^{-3} w_0(R) \le w(y) \le \beta^3\, w_0(R)$. Hence

$$\mu(B^*(x,R)\setminus B_{R/2}) \le \beta^3\, w_0(R)\,|B^*(x,R)| \le e\beta^3\, w_0(R)\,|B_R|.$$

On the other hand note that if $y \in B(x,R)\setminus B_{R/2}$ then $R/2 \le |y| \le 4R$ and consequently $\beta^{-2} w_0(R) \le w(y) \le \beta^2\, w_0(R)$. This yields

$$\begin{aligned} \mu(B(x,R)) &\ge \mu(B(x,R)\setminus B_{R/2}) \ge \frac{1}{\beta^2}\, w_0(R)\,|B(x,R)\setminus B_{R/2}| \\ &\ge \frac{1}{2\beta^2}\, w_0(R)\,|B_R|. \end{aligned}$$

□

Remark. Observe that the only thing that we have really proved here is that a uniform weakly doubling property implies a uniform micro-doubling property. The reciprocal is not true. In the last Section, we show an example of a density for which the associated measure is uniformly n-micro-doubling but is not uniformly weakly doubling.

3.4. **Decreasing densities.**

Radial measures with decreasing densities have some interesting and sometimes surprising properties. For instance, to ensure that a measure with such a density is uniformly weakly bounded, it is enough to check this condition just in one concrete Euclidean space.

Proposition 3.13. *Let w_0 be a decreasing function over $[0,\infty)$. If μ is weakly doubling in $\mathbb{R}^N$ for some $N \in \mathbb{N}$, then μ is uniformly weakly doubling in $\mathbb{R}^n$ for all $n \ge N$.*

Proof. Assume that μ is weakly doubling with constant K_1 in $\mathbb{R}^N$ for some $N \in \mathbb{N}$. By Theorem 3.3 we only need to show that w is essentially constant over dyadic annuli. Given $R > 0$, if $R \le |x| \le 2R$, since w_0 is decreasing, one has $w_0(2R) \le w(x) \le w_0(R)$. Thus, we only need to check that $w(R) \le Cw(2R)$ with C independent of R. To do so, consider a point $z \in \mathbb{R}^N$ so that $|z| = R/2$. Take $z' = 3z$ and $z'' = 5z$ and consider the balls $B = B(z,R/2)$, $B' = B(z',R/2)$ and $B'' = B(z'',R/2)$.



By the weakly doubling property of μ we have that $\mu(B) \leq K_1\mu(B') \leq K_1^2\mu(B'')$. For all $y \in B$ we have $w_0(R) \leq w(y)$ and for all $y \in B''$ we have $w(y) \leq w_0(2R)$, this yields

$$w_0(R) \leq \frac{\mu(B)}{|B|} \leq K_1^2\,\frac{\mu(B'')}{|B''|} \leq K_1^2\,w_0(2R).$$

□

Also, if the density is decreasing, the reciprocal implication in Theorem 3.1 is true.

Proposition 3.14. *Let w_0 be a decreasing function over $[0,\infty)$. Assume that there exists $C > 0$ so that for some $N > 0$ one has $\|M_\mu\|_{L^1_{\text{rad}}(\mathbb{R}^n,d\mu)\to L^{1,\infty}_{\text{rad}}(\mathbb{R}^n,d\mu)} \leq C$ for all $n \geq N$. Then μ is uniformly weakly doubling in $\mathbb{R}^n$ for all $n \geq N$.*

Observe that by Theorem 3.2, the hypothesis in the previous proposition also implies that M_μ is uniformly bounded on $L^p(\mathbb{R}^n, d\mu)$ for all $n \geq N$. It is remarkable that a boundedness condition over radial functions implies one for general functions. The proof of this Proposition is based on differentiation through dimensions.

Proof. Assume that M_μ satisfies the weak $L^1(\mu)$ inequality

$$\mu\left(\{x \in \mathbb{R}^n : M_\mu f(x) > \lambda\}\right) \leq \frac{C_*}{\lambda}\int_{\mathbb{R}^n} |f(x)|\,d\mu(x),$$

for all $f \in L^1_{rad}(\mathbb{R}^n)$ and $n \geq N$ for certain $N \in \mathbb{N}$, with C_* independent of f and n. As before, it is enough to show that w is essentially constant over dyadic annuli. Since w_0 is decreasing, it suffices to prove that $w_0(R) \leq Cw_0(2R)$ with C independent of R. Fix $R > 0$ and for each $n \geq N$, take $z^n \in \mathbb{R}^n$ so that $|z^n| = R$. Set $f = \chi_{B_\varepsilon}$ with $0 < \varepsilon < R$ and

$$\lambda = \frac{\mu(B(z^n, R+\varepsilon))}{\mu(B_\varepsilon)}.$$

Clearly we have $B_R \subset \{x \in \mathbb{R}^n :\ M_\mu f(x) > \lambda\}$, which together with our hypothesis gives

$$\mu(B_R) \leq \frac{C_*}{\lambda}\int_{\mathbb{R}^n} |f(x)|\,d\mu(x) = C_*\,\mu(B(z^n, R+\varepsilon)).$$

Letting $\varepsilon \to 0^+$, by monotonicity we obtain $\mu(B_R) \leq C_*\mu(B(z^n,R))$. In particular we have

$$\frac{\mu(B_R)}{|B_R|} \leq C_*\frac{\mu(B(z^n,R))}{|B(z^n,R)|},$$

and now we differentiate through dimensions. Letting $n \to \infty$ in the last inequality, Lemma 3.11 gives $w_0(R) \leq C_*w_0(\sqrt{2}R)$ from which we deduce that $w_0(R) \leq Cw_0(2R)$ with $C = C_*^2$. □

We now make a connection between the previous hypotheses and the Muckenhoupt A_1 condition. We recall that for a positive $w \in L^1_{\text{loc}}(\mathbb{R}^n)$ the A_1-constant we call $[w]_{A_1(\mathbb{R}^n)}$ is defined as the smallest constant C so that

$$\fint_{B(x,R)} w(y)\,dy \le C \operatorname*{ess\,inf}_{y \in B(x,R)} w(y),$$

for all $x \in \mathbb{R}^n$ and $R > 0$. Under the hypothesis that w_0 is decreasing in $[0,\infty)$ and $d\mu(x) = w_0(|x|)\,dx$ is uniformly weakly doubling we will show that $w(x) = w_0(|x|) \in A_1(\mathbb{R}^n)$ for all sufficiently large n. Moreover, the A_1-constants can be bounded independently of n. The precise statement is the following.

Proposition 3.15. *Let w_0 be a decreasing function over $[0,\infty)$ and essentially constant over dyadic intervals with constant β. Then there exist $N > 0$ and $C > 0$ only depending on β so that $[w]_{A_1(\mathbb{R}^n)} \le C$ for all $n \ge N$.*

Proof. Assume that w is essentially constant over dyadic intervals with constant β and take $n \ge N$ for the N given in Lemma 3.9. We have to find a constant $C > 0$ so that for all $x \in \mathbb{R}^n$ and $R > 0$ one has

$$\frac{\mu(B(x,R))}{|B(x,R)|} \le C \inf_{y \in B(x,R)} w(y).$$

If $|x| \le 2R$, since w is decreasing and using Lemma 3.9 and that w is essentially constant over dyadic annuli one has

$$\frac{\mu(B(x,R))}{|B(x,R)|} \le \frac{\mu(B_R)}{|B_R|} \le 2\beta\, w_0(R) \le 2\beta^3\, w_0(4R).$$

This finishes the proof in this case because now using again that w_0 is decreasing we obtain

$$w_0(4R) \le \inf_{y \in B_{4R}} w(y) \le \inf_{y \in B(x,R)} w(y).$$

If $|x| \ge 2R$ then w is essentially constant over $B(x,R)$. Indeed if $y \in B(x,R)$ then $|x|/2 \le |y| \le 3|x|/2$, which implies that $\beta^{-1} w(x) \le w(y) \le \beta\, w(x)$. Hence,

$$\frac{\mu(B(x,R))}{|B(x,R)|} \le \beta\, w(x) \le \beta^2 \inf_{y \in B(x,R)} w(y).$$

□

The amazing thing here is that the reciprocal statement of this last proposition is also true.

Proposition 3.16. *Let w_0 be a non-negative function over $[0,\infty)$ and set $w(x) = w_0(|x|)$ so that $w \in A_1(\mathbb{R}^n)$ for $n \ge N > 0$. If there exists $C_* > 0$ so that $[w]_{A_1(\mathbb{R}^n)} \le C_*$ for all $n \ge N$, then w_0 is essentially constant over dyadic intervals and comparable with a decreasing function.*

Proof of Proposition 3.16. It is enough to prove that w is comparable with a decreasing function $\tilde{w}$ because then $\tilde{w} \in A_1$ and, therefore, $\tilde{w}$ is (weakly) doubling. By Proposition 3.13 and Theorem 3.3 we would have that $\tilde{w}$ is essentially constant over dyadic intervals, and hence so is w.

It suffices to check that there is a constant $q > 0$ so that $w_0(t) \le q\, w_0(s)$ whenever $0 \le s \le t$. For $0 < s < t$ and $n \ge N$, take $x_n, y_n \in \mathbb{R}^n$ so that x_n is a multiple of y_n and $|x_n| = s$, $|y_n| = t$. Consider the ball $B(x_n, R)$ with $R^2 = t^2 - s^2$.



By hypothesis we have, in the almost everywhere sense,

$$\frac{\mu(B(x_n,R))}{|B(x_n,R)|} \leq C_* \, w(x_n) = C_* \, w_0(s).$$

Letting $n \to \infty$, by the differentiation through dimensions of Lemma 3.11 we get that

$$w_0(t) = w_0(\sqrt{s^2+R^2}) \leq C_* \, w_0(s),$$

for almost every $s \leq t$. □

4. FURTHER RESULTS AND REMARKS

4.1. **Examples of measures that are not uniformly weakly-doubling.** Each measure μ with density $w(x) = |1-|x||^{-\alpha}$ for $0 < \alpha < 1$ is doubling on $\mathbb{R}^n$ for each $n \in \mathbb{N}$, but is not uniformly weakly doubling. To see this observe that in each $\mathbb{R}^n$ one has $\mu(B_1) \geq C_\alpha n^\alpha |B_1|$. Using differentiation through dimensions, if $|z| = 1$, then $\mu(B(z,1))/|B(z,1)| \overset{n\to\infty}{\longrightarrow} w(\sqrt{2}) = (\sqrt{2}-1)^{-\alpha}$. This says that the measures of the intersecting balls B_1 and $B(z,1)$ are not comparable with constants independent of the dimension.

Using this idea we can also prove that there are not uniform weak $L^1(\mu)$ bounds for the associated maximal operator.

Theorem 4.1. *Let μ be the measure defined above and let $C_{1,n}$ be the smallest constant satisfying*

$$\mu(\{x \in \mathbb{R}^n : M_\mu f(x) > \lambda\}) \leq \frac{C_{1,n}}{\lambda} \|f\|_{L^1(\mathbb{R}^n,\mu)},$$

for all $f \in L^1(\mathbb{R}^n,\mu)$. Then, one has $C_{1,n} \geq cn^\alpha$.

Proof. Using discretization (see [19] and [24]), or the argument in the proof of Proposition 3.14, we may consider M_μ acting over finite sums of Dirac deltas instead of integrable functions. For $\delta_0 = \lim_{\varepsilon\to 0} \chi_{B_\varepsilon}(x)/\mu(B_\varepsilon)$, in the sense of distributions, the weak $L^1(\mu)$ inequality reads

$$\mu(\{x \in \mathbb{R}^n : M_\mu \delta_0(x) > \lambda\}) \leq \frac{C_{1,n}}{\lambda}. \tag{4.1}$$

Note that $M_\mu\delta_0(x) = 1/\mu(B(x,|x|))$ for each $x \neq 0$, which makes $M_\mu\delta_0$ a radially decreasing function. Then taking $\lambda = M_\mu\delta_0(z)$ with $|z| = 1$ we have $\{x \in \mathbb{R}^n : M_\mu\delta_0(x) > M_\mu\delta_0(z)\} \subset B_1$ and from (4.1) we obtain

$$C_{1,n} \geq \frac{\mu(B_1)}{\mu(B(z,1))}.$$

This, together with the previous observation that $\mu(B_1) \geq Cn^{\alpha}|B_1|$ and that $\mu(B(z,1))/|B_1| \to w(\sqrt{2})$ as $n \to \infty$, proves the result. $\square$

Remark. One can prove with some extra work that μ is uniformly n-micro-doubling. It is also weakly doubling but in this case with a constant $K_1 \sim n^{\alpha}$ in each $\mathbb{R}^n$. In particular, Theorems 1.1 and 4.1 imply that $C_{1,n} \leq c\, n^{1+\alpha} \log n$.

4.2. **Families of measures changing with the dimension.** As we have seen in the preceding sections, maximal operators associated with measures given by power densities have L^p operator norms bounded with respect to the dimension. An interesting observation by J.M. Aldaz and J. Pérez Lázaro in [3] showed that given an exponent p (as large as wanted) there exist families of power weights such that the L^p bounds of the associated maximal operators grow to infinity as $n \longrightarrow \infty$. The twist here is that the powers change from one dimension to another. To be more precise they considered measures $\nu_{\alpha,n}$ given by the densities $|x|^{-\alpha n}$ over $\mathbb{R}^n$ with $0 < \alpha < 1$. Their result is the following (see Theorem 3.12 in [3]).

Theorem 4.2. *Given $p_0 \in [1,\infty)$, there exist $\alpha_0 \in (0,1)$ and $a > 1$ such that for all $p \in [1,p_0]$ and all $\alpha \in [\alpha_0, 1)$ one has*

$$c_{\nu_{\alpha,n},p} \geq \frac{a^{(1-\alpha)n}}{6}.$$

It is implicit in the proof given in [3] that $\alpha_0 \longrightarrow 1$ as $p_0 \longrightarrow \infty$. This leads to the question of whether, fixing α, $M_{\nu_{\alpha,n}}$ may satisfy a uniform L^p bound for large p. We can apply the method used in Theorem [15] for the Gaussian measure to show that this is not the case when $\alpha > 1/2$.

Theorem 4.3. *For each $\alpha \in (1/2,1)$ there exists a constant $a > 1$ such that for all $p \in [1,\infty)$*

$$c_{\nu_{\alpha,n},p} \geq ca^{n/p},$$

even if the action is restricted to radially decreasing functions.

The proof of this result can be found in [14] and in [4].

A consequence of this result is that for these families of measures the constants of the n-micro-doubling and the weak doubling conditions grow to infinity with the dimension (see Theorems 1.1, 3.1 and 3.6).

REFERENCES

[1] J.M. Aldaz, *Dimension dependency of the weak type (1,1) bounds for maximal functions associated to finite radial measures.* Bull. Lond. Math. Soc. **39** (2007), 203–208.

[2] J.M. Aldaz, *The weak type (1,1) bounds for the maximal function associated to cubes grow to infinity with the dimension.* Ann. of Math. (2) **173** (2011), no. 2, 1013-1023.

[3] J.M. Aldaz, J. Pérez Lázaro, *Dimension dependency of L^p bounds for maximal functions associated to radial measures.* Positivity **15** (2011), 199–213.

[4] J.M. Aldaz, J. Pérez Lázaro, *On high dimensional maximal operators.* Banach J. Math. Anal. **7** (2013), no. 2, 225–243.

[5] J.M. Aldaz, J. Pérez Lázaro, *The best constant for the centered maximal operator on radial decreasing functions.* Math. Inequal. Appl. **14** (2011), no. 1, 173–179.

[6] G. Aubrun, *Maximal inequality for high-dimensional cubes.* Confluentes Math. **1** (2009), no. 2, 169–179.

[7] J. Bourgain, *On high-dimensional maximal function associated to convex bodies.* Amer. J. Math. **108** (1986), no. 6, 1467–1476.

[8] J. Bourgain, *On the L^p-bounds for maximal functions associated to convex bodies in R^n*. Israel J. Math. **54** (1986), no. 3, 257–265.
[9] J. Bourgain, *On dimension free maximal inequalities for convex symmetric bodies in $\mathbb{R}^n$*. Geometrical aspects of functional analysis (1985/86), 168–176, Lecture Notes in Math., **1267**, Springer, Berlin, 1987.
[10] J. Bourgain, *Averages in the plane over convex curves and maximal operators.* J. Analyse Math. **47** (1986), 69–85.
[11] J. Bourgain, *On the Hardy-Littlewood maximal function for the cube* arXiv:1212.266v1.
[12] A. Carbery, *An almost-orthogonality principle with applications to maximal functions associated to convex bodies.* Bull. Amer. Math. Soc. (N.S.) **14** (1986), no. 2, 269–273.
[13] A. Criado, *On the lack of dimension free estimates in L^p for maximal functions associated to radial measures.* Proc. Roy. Soc. Edinburgh Sect. A **140** (2010), no. 3, 541-552.
[14] A. Criado, *Problems of harmonic analysis in high dimensions.* Ph.D. thesis, Universidad Autónoma de Madrid, 2012.
[15] A. Criado, P. Sjögren, *Weak L^p bounds for maximal functions associated to rotational invariant measures in $\mathbb{R}^n$ for large n.* To appear in J. Geom. Anal.
[16] J. Duoandikoetxea, L. Vega, *Spherical means and weighted inequalities.* J. London Math. Soc. (2) **53** (1996), no. 2, 343–353.
[17] J. García-Cuerva, J.L. Rubio de Francia, *Weighted norm inequalities and related topics.* North-Holland Mathematics Studies, 116. Notas de Matemática [Mathematical Notes], 104. North-Holland Publishing Co., Amsterdam, 1985.
[18] A.M. Garsia, *Topics in almost everywhere convergence.* Lectures in Advanced Mathematics, 4 Markham Publishing Co., Chicago, Ill. 1970.
[19] M. de Guzmán, *Differentiation of Integrals in $\mathbb{R}^n$.* Lecture Notes in Math., Vol. 481. Springer-Verlag, Berlin-New York, 1975.
[20] A. Infante, *Análisis armónico para medidas no doblantes: operadores maximales con respecto a la medida gaussiana*, Ph.D. thesis, Universidad Autónoma de Madrid, 2005.
[21] A. Infante, *Free-dimensional boundedness of the maximal operator.* Bol. Soc. Mat. Mexicana (3) **14** (2008), no. 1, 67–73.
[22] E. Lindenstrauss, *Pointwise theorems for amenable groups.* Invent. Math. **146** (2001), no. 2, 259–295.
[23] M.T. Menárguez, *Técnicas de discretización en Análisis Armónico para el estudio de acotaciones débiles de operadores maximales e integrales singulares.* Ph. D. thesis, Universidad Complutense, Madrid, 1990.
[24] M.T. Menárguez, F. Soria, *Weak type (1,1) inequalities for maximal convolution operators.* Rend. Circ. Mat. Palermo (2) **41** (1992), no. 3, 342–352.
[25] M.T. Menárguez, F. Soria, *On the maximal operator associated to a convex body in $\mathbb{R}^n$.* Collect. Math. **3** (1992), 243–251.
[26] B. Muckenhoupt, E.M. Stein, *Classical expansions and their relation to conjugate harmonic functions.* Trans. Amer. Math. Soc. **118** 1965 17–92.
[27] D. Müller, *A geometric bound for maximal functions associated to convex bodies.* Pacific J. Math. **142** (1990), no. 2, 297–312.
[28] A. Naor, T. Tao *Random martingales and localization of maximal inequalities*, J. Funct. Anal. **259** (2010), no. 3, 731-779.
[29] J.L. Rubio de Francia, *Maximal functions and Fourier transforms.* Duke Math. J. **53** (1986), no. 2, 395-404.
[30] E.M. Stein, *Maximal functions. I. Spherical means* Proc. Nat. Acad. Sci. U.S.A. **73** (1976), no. 7, 2174–2175.
[31] E.M. Stein, *The development of square functions in the work of A. Zygmund.* Bull. Amer. Math. Soc. (N.S.) **7** (1982), no. 2, 359–376.
[32] E.M. Stein, J.O. Strömberg, *Behavior of maximal functions in R^n for large n.* Ark. Mat. **21** (1983), no. 2, 250–269.
[33] E.M. Stein, G. Weiss, *Introduction to Fourier analysis on Euclidean spaces.* Princ. Math. Ser., No. 32. Princeton University Press, Princeton, N.J., 1971.

ALBERTO CRIADO
DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN
E-mail address: `alberto.criado@uam.es`

FERNANDO SORIA
DEPARTAMENTO DE MATEMÁTICAS AND INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UCM-UC3M, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID, SPAIN
E-mail address: `fernando.soria@uam.es`